



Clinicopathological profile of hyperpigmented skin lesions- A prospective study

Dr Sirish Aryasomayajula MD¹, Dr Manimala Danda MD¹, Varun Kamidi MD¹, Aswini Gude MD², Padmapriya Mamillapalli³, Kirankumar Epari MD⁴, Vahini Gudeli MD⁵, Dr. B. A. Ramakrishna MD⁵

Abstract: **Introduction:** Hyperpigmentation is one of the most common reaction to inflammatory, benign and malignant lesions of the skin. These disorders comprise heterogeneous group of diseases of epidermal and dermal hyperpigmentation divided into various types according to etiology and pathology. Correct diagnosis of these hyperpigmented lesions is linked to histopathologic examination of skin biopsies with clinical correlation. **Aim:** To study the spectrum of hyperpigmented skin lesions with reference to age and sex distribution. To study the histopathology of hyperpigmented skin lesions in correlation with clinical features. **Materials and Methods:** This prospective cohort study was conducted at Department of Pathology, at Alluri Sitaramaraju Academy of Medical Sciences, Eluru, Andhra Pradesh, India, which included 80 patients who were clinically diagnosed with hyperpigmented skin lesions in all age groups from July 2014 to August 2016. Frequency and percentage statistics was used to present the results. **Results:** Out of 80 cases, 34 cases of inflammatory lesions, 23 cases of benign lesions and 23 cases of malignant lesions were reported. Among the post inflammatory lesions the majority were classical Lichen planus. **Conclusion:** Most common lesion was lichen planus and its variants with highest incidence in females and age group greater than 60 years. Histopathological diagnosis with clinical correlation aids in effective management of the patients.

Keywords: Hyperpigmentation, Lichen planus, Becker's nevus, Spindle cell Nevus of Reed, Spitz nevus.

AFFILIATIONS

¹Assistant Professor, Department of Pathology, Gayatri Vidhya Parishad Institute of Healthcare and Medical Technology, Visakhapatnam, Andhra Pradesh, India

²Associate Professor, Department of Pathology, Gayatri Vidhya Parishad Institute of Healthcare and Medical Technology.³

³MBBS, DDVL Consultant Dermatologist

⁴Professor, Department of Pathology, Gayatri Vidhya Parishad Institute of Healthcare and Medical Technology, Visakhapatnam, Andhra Pradesh, India

⁵Professor, Department of Pathology, Alluri Sita Ramaraju Academy of medical sciences, Eluru, Andhra Pradesh, India

CORRESPONDING AUTHOR

Dr. Manimala Danda

INTRODUCTION

Pigmented skin lesions are one of the most frequent problem encountered by the dermatologist, ranging from hypopigmentation to hyperpigmentation. Most of them are benign, a small percentage of cases are malignancies ^[1]. Disorders of pigmentation can result from migration abnormalities of melanocytes from neural crest to the skin during embryogenesis, improper transfer of melanosomes to keratinocytes and variation in melanin synthesis ^[2].

Hyperpigmentation is one of the most common reaction to inflammatory, benign and malignant lesions of the skin. These disorders comprise heterogeneous group of diseases of epidermal and dermal hyperpigmentation divided into various types according to aetiology and pathology ^[3].

Correct diagnosis of these hyperpigmented lesions is often linked to histopathologic examination of skin biopsies with clinical correlation, as there is evolution of skin lesions into different stages in disease progression ^[3]. The significance of precise diagnosis is emphasized in hyperpigmentary skin lesions as the underlying diseases have varying aetiologies for the purpose of treatment ^[2].

The present study was conducted to understand the spectrum of hyperpigmented skin lesions with reference to age, sex and to describe the utility of histology in aid with clinical correlation for targeted and more effective therapy.

MATERIALS AND METHODS

The present study is a two year prospective study from July 2014 to August 2016, included 80 patients who were clinically diagnosed with hyperpigmented skin lesions in all age groups from department of Dermatology, Surgery and Plastic surgery at Alluri Sita Ramaraju Academy of Medical Sciences, Eluru.

Pertinent clinical details like age, duration of the lesion, site of the lesion, significant medical history were taken and entered in the proforma. The lesions were biopsied after taking consent from the patient. Punch biopsy of 4mm was done for smaller lesions while larger lesions were excised and sent in 10% formaldehyde solution. Tissue was processed and stained with routine Haematoxylin and Eosin stain for histopathological examination. Special stains like Masson Fontana and Melanin bleach were performed whenever required. The results were analyzed using descriptive studies.

Inclusion Criteria

Non-neoplastic and neoplastic pigmented skin lesions in various age groups, received in the department of pathology were included in the study.

Exclusion Criteria

1. All hypopigmented skin lesions and inadequate biopsies.
2. Post chemotherapy and post-radiotherapy skin lesions.
3. Unsuitable for biopsy due to ulceration or infection

RESULTS

The period of present study was of 2 years duration from July 2014 to August 2016. Total number of biopsies included were 80. These biopsies after histopathological evaluation were categorized under inflammatory, benign, and malignant lesions. The present study includes 34 cases (42.5%) of inflammatory lesions, 23 cases (28.75%) of benign lesions and 23 cases (28.75%) of malignant lesions. Among the post inflammatory lesions the highest incidence were classical Lichen planus, which were 15 cases out of 34 cases i.e. 44.12% (Table 1).

In the present study 23 cases of benign lesions were noted, 14 were diagnosed as Nevus (60.8%) followed by 9 cases of Seborrheic keratosis (39.13%) (Table 1). 23 malignant lesions were included, majority were Basal cell carcinoma (16 cases 69.56%), out of which four were diagnosed as Pigmented variant of BCC, followed by 7 cases of malignant melanoma (30.4%) (Table 1). Overall the incidence of cases were in fifth decade followed by fourth and second accounting for 48.75%, 13.75% and 12.5% respectively. Male preponderance was seen (Table 2). In the present study, positive correlation was seen between clinical diagnosis and histopathological diagnosis in 90 % of cases. In 10% cases discordance was noted and final diagnosis was given based on histopathological findings (Table 3).

DISCUSSION

In the present study of hyperpigmented skin lesions the post inflammatory hyperpigmentation is the most common sequelae among the inflammatory dermatosis. It is an acquired hypermelanosis occurring after cutaneous inflammation or injury that can arise in all skin types [4]. This study analyzed the spectrum of hyperpigmented skin lesion and ascertain the significance of histopathological diagnosis for its accurate characterization. Most common age group involved in post inflammatory cases was sixth decade followed by third and fifth accounting for 44.12%, 17.65% and 14.70% of cases respectively. Females were more commonly affected than males. As per Daniel L. Stulberg et al [5] and Nicole Clark et al [5] post inflammatory hyper pigmentation can occur at any age with increase in incidence between 15- 60 years affecting both males and females in an equal proportion affecting the extremities and face. Female preponderance was noted in the studies done by Sontheimer RD et al [6].

Majority of cases in the present study were classical lichen planus and its variants with equal sex distribution and age group >50 years were most commonly affected which correlated with the results of Mrutyunjayappa et al [3] and sneha et al [2].

In the present study clinically lichen planus and its variants presented as hyperpigmented, pruritic, flat, papules and plaques affecting all over the body and on histopathological examination showed hyperkeratosis, acanthosis,

hypergranulosis, basal cell degeneration, saw too thing of rete ridges, band likely lymphohistiocytic infiltrate at dermo epidermal junction

(Figure 1) which correlated with the histological features in the studies of Ellis FA et al^[7], Sontheimer RD et al^[6] and Sehgal VN et al^[8].

In the present study 3 cases of lichen simplex chronicus were reported which presented as multiple itchy plaques over back, upper limb and lower limb. Histopathological examination revealed hyperkeratosis, parakeratosis, acanthosis, irregular elongation of rete ridges, wedge shaped hypergranulosis, broadening of dermal papillae, superficial perivascular infiltrate. Similar findings were noted in the study of Lotti Tetal^[9] and Rajasekhar Netal^[10].

In the present study two cases of Prurigonodularis were reported involving the extremities as multiple hyperpigmented papules and plaques. Histologically lesion showed pronounced hyperkeratosis, irregular acanthosis, papillomatosis, irregular downward proliferation of epidermis, and lymphocytic infiltrate in dermis, vertically oriented collagen bundle correlated with the features of Lee RM et al^[11], Taefeh norooz H et al^[12] and Thadeus J^[13].

2 cases of morphea were diagnosed which presented as hyperpigmented multiple papules over extremities and abdomen. On histopathological examination showed mild lymphocytic infiltrate and thickened collagen bundles in reticular dermis. Jaworsky C et al^[14] and Dhar S et al^[15] also found similar results to our study.

Among the benign lesions Seborrheic keratosis is the most common lesion which develops from the proliferation of keratinocytes of the epidermis. In the present study we got 9 cases of seborrheic keratosis out of which 7 were pigmented, 2 cases were reported on the breast, left side of the mandible, each one case in the lumbar region, lower lip, scalp, forehead and thigh. Males were more commonly affected with most common age group involved was sixth decade.

As per study conducted by Arthur K Balin et al^[16] and Tindall et al^[17] seborrheic keratosis is the most common benign tumor in older individuals. Pigmented seborrheic keratosis is easily mistaken for malignant melanoma and in the present study provisional clinical diagnosis in one case was kept as melanoma. Based on histological picture seborrheic keratosis was diagnosed.

In the present study 2 cases of Becker's Nevi were reported and both were males and had the lesion since birth on forearm which gradually grew and was associated with hypertrichosis after puberty (Figure 2). No other anomalies such as hypoplasia of underlying structures were noted in the present study.

In the present study a single case of Spindle cell nevus of Reed was reported in a 18 year old female who presented with uniformly pigmented plaque over the legs. Histopathological examination revealed monomorphic spindled epithelioid melanocytes arranged in nests within the epidermis and papillary dermis with occasional upward inter epidermal progression of melanocytes. The findings in our study correlated with the study conducted by K.K. Boneti et al^[18].

In the present study two cases of Nevus sebaceous presented as a circumscribed lesion in a 19 year old male and 47 years female over the neck and infrascapular region. The commonest benign and malignant tumour arising in nevus sebaceous is Syringocystadenoma papilliferum and Basal cell carcinoma respectively.

A single case of Spitz nevus was reported in a 12 year old male who presented with hyperpigmented lesion over the hand. According to Wettengel G.V et al^[19] Spitz nevus is a proliferation of melanocytes can be diagnosed wrongly as malignant melanoma and vice versa. Clinically in our study the lesion was diagnosed accurately as Nevus. In our case histopathology did not show any neoplastic changes.

Among the 23 malignant lesions, Basal cell carcinoma (BCC) accounted for 16 cases accounting for 69.56% cases. Out of 16 cases, 4 were pigmented variant of BCC. Majority of cases in the present study were seen in the 6th decade with female preponderance which correlated with the results of Saraswathy Sreeram et al^[20].

In the present study 7 cases of Malignant Melanoma were reported accounting for 30.43% of all malignant lesions. The maximum numbers of cases were seen in the 6th decade with female preponderance which correlated with features of

Wanebo *et al* [21]. In the diagnosis of hyperpigmentary skin lesions morphology, distribution, pattern and extent of lesions are helpful to make an accurate provisional clinical diagnosis and study of skin biopsy gives a confirmatory diagnosis. In the present study, positive correlation was seen between clinical diagnosis and histopathological diagnosis in only 90 % of cases thus emphasizing the importance and utility of histopathology in arriving at a conclusive diagnosis.

Limitation

The limitation of the study was size of the sample was small and there was no follow up of the cases.

TABLES

Type of lesion	Number of cases
Inflammatory lesions	34(100%)
Classical Lichen planus	15 (44.12%)
Lichenoid reactions	7(20.5%)
Lichen simplex chronicus	3(8.8%)
Prurigonodularis	2(5.8%)
Pruritic papular eruption	2(5.8%)
Morphea	2(5.8%)
Fixed drug eruption	1(2.9%)
Urticariapigmentosa	1(2.9%)
Incontinentiapigmenti	1(2.9%)
Benign lesions	24(100%)
Seborrheickeratosis	9(39.13%)
CompoundNevus	3(13.04%)
Junctional melanocytic Nevus	2(8.7%)
Beckers Nevus	2(8.7%)
Congenital melanocytic Nevus	2(8.7%)
Nevus sebaceous	2(8.7%)
Epidermal Nevus	2(8.7%)
Spindle cell Nevus of Reed	1(4.35%)
SpitzNevus	1(4.35%)
Malignant lesions	23(100%)
Basal cell carcinoma	16(69.56%)
Malignant melanoma	7(30.4%)

Table 1- Showing the distribution of skin lesions

Age in years	Male	Female	Total	Percentage
Birth - 10	0	3	3	3.75%
11-20	5	3	8	10%
21-30	6	4	10	12.50%
31-40	4	5	9	11.25%
41-50	8	3	11	13.75%
>50	19	20	39	48.75%
Total	41	39	80	100%

Table 2 - Age &sex distribution of hyperpigmented skin lesions

Clinical diagnosis	Histopathological diagnosis
Fixed drug eruption	Pruritic papular eruption
Eczema	Urticariapigmentosa
Pellagra	Incontinentia pigment
Prurigo simplex	Post inflammatory hyper pigmentation
Discoid lupus erythematosus	Pruritic papular eruption
Malignant melanoma	Seborrheic keratosis

Table 3 - Cases diagnosed on histopathology with variation to clinical diagnosis

FIGURES




Figure1:Lichen planus-Show structure of epidermis with hyper keratosis and hyper granulos is with collection of band like lymphocytes within the dermis.



Figure 2: - Becker's Nevus - Pigmented irregular warty growth with hypertrichosis

CONCLUSION

Lichen planus and its variants were the most common hyperpigmented skin lesions encountered in this study. As there is a significant overlap in clinical presentation, specific diagnosis of hyperpigmented lesions were based on histopathologic findings and its clinical correlation for definite treatment. Histopathological diagnosis with clinical correlation aids in the effective management of patients.

REFERENCES

1. DowerahSwagata, Deori Rashmi. A histopathological study of pigmented lesions of skin: melanoma and its clinical mimics.GJRA.vol-6, issue 11 Nov; pg284-285.
2. SnehaRavindran, Kalla Ravi Teja, Arun Kumar S, Jonathan Arnold, Balaji S. Histopathological and Clinical Correlation of Hyperpigmented Skin Lesions.Sch J App Med Sci, Nov, 2020; 8(11):2472-2477.
3. Mruthyunjayappa S, Mahantappa H, Gopal M G, Venugopal SB. A study of spectrum of histopathological features in

- patients presenting with hyperpigmented skin lesions. Arch Med Health Sci. 2016; 4: p. 189-95.
5. Erica C. Davis, Valarie D. Callender, Post inflammatory hyperpigmentation a review of the epidemiology, clinical features, and treatment options in skin of color. The journal of clinical and Aesthetic dermatology.vol3, July 2010,page20-31.
 6. Daniel L. Stulberg, and Nicole Clark,M. Common Hyperpigmentation disorders in adults, Postinflammatoryhyperpigmentation, Melanoma, Seborrheic Keratosis, Acanthosis, Nigricans, M elasma, Diabetic dermopathy, Tinea Versicolor and Post inflammatory hyperpigmentation- American family Physician vol 68, 10/2003 Nov15
 7. Sontheimer RD. Lichenoid tissue reaction/interface dermatitis: Clinicaland Histological perspectives. J Invest Dermatol. 2009; 129: p.1088-99.
 8. Ellis FA. Histopathology of lichen planus based on analysis of one hundredbiopsy specimens. The journal of investigative dermatology. 1967; 48(2):143-148.
 9. Sehgal VN, Srivastava G, Sharma S, Sehgal S, Verma P. Lichenoid tissue reaction/interface dermatitis: Recognition, classification, etiology and clinicopathological overtones. Indian Journal of Dermatol venereal Leprol 2011; 77(4):418-430.
 10. Lotti T, Buggiani G, Prignano F. Prurigo Nodularis and Lichen simplex chronicus.
 11. Dermatology Therapy 2008; 21: 42-46.
 12. Rajasekhar N, Thippeswamy C, Prasanna NB. Lichen simplex chronicus of scrotum. Indian Journal of Dermatol venereal Leprol 1999; 65:91-92.
 13. Lee RM, Shumack S. Prurigo Nodularis: A review. Australian Journal of Dermatology 2005; 46:211-220.
 14. Taefehnorooz H, Truchetet F, Barbaud A, Schmutz J, Bursztejn A. Efficacy of Thalidomide in the treatment of Prurigo Nodularis. Acta Derma-Venereol 2011;91
 15. Thadeus J, Thappa DM, Ratnakar C. Prurigo Nodularis with Cutaneous hom. Ind J Dermatol 1997; 42(3):188-190.
 16. Jaworsky C, Winfield H. Connective tissue diseases. Lever's Histopathology of skin. Tenth edition. Published by Wolterskluwer 2009:296-298.
 17. Dhar S, Dhar S. Why lesions of Morphea are often hyperpigmented? Indian Journal of Dermatol venereal Leprol 1996; 62(2):131.
 18. Arthur K Balin . The Sally Balin Medical Centre. Seborrheic Keratosis- Medscape 17. Tindall JP. Smith JG, Jr. Skin lesion of the aged and their association with internal changes. JAMA 1963 Dec 21. 186:1039-42 [Medlin
 19. K. Boneti, J. Piñeiro-Maceira, F. B. Pereira and C. B. Barcaui, "Reed Nevus (Pigmented Spindle-Cell Nevus): A Report of Three Cases with Distinct Dermoscopic Patterns," Anais Brasileiros de Dermatologia, Vol. 85, No. 4, 2010, pp.531
 20. Wettengel G. V., Draeger J., Kiesewetter F., Schell H., Neubauer S., Gebhart E., Differentiation between Spitz nevi and malignant melanomas by interphase fluorescence in situ hybridization, Int J Oncol, 1999. 14(6):1177-1183.
 21. Saraswathy Sreeram, Flora Dorothy Lobo, Ramdas Naik, Urmila Niranjana Khadilkar, Hema Kini, Ullal Anand Kini; Morphological Spectrum of Basal Cell Carcinoma in Southern Karnataka: Journal of Clinical and Diagnostic Research. 2016 Jun, Vol-10(6):EC04-EC07.
 22. Wanebo HJ, Cooper PH, Young DV. Prognostic factors in head and neck melanoma. Effect of lesion location. Cancer 1988; 62:831-7